

SHORT REPORT



Low prevalence of anemia among Shuar communities of Amazonian Ecuador

Alicia M. DeLouize¹ | Melissa A. Liebert² | Felicia C. Madimenos^{3,4} | Samuel S. Urlacher^{5,6} | Joshua M. Schrock¹ | Tara J. Cepon-Robins⁷ | Theresa E. Gildner⁸ | Aaron D. Blackwell⁹ | Christopher J. Harrington¹ | Dorsa Amir¹⁰ | Richard G. Bribiescas¹¹ | James Josh Snodgrass^{1,12} | Lawrence S. Sugiyama¹

¹Department of Anthropology, University of Oregon, Eugene, Oregon

²Department of Anthropology, Northern Arizona University, Flagstaff, Arizona

³Department of Anthropology, Queens College (CUNY), Flushing, New York

⁴New York Consortium in Evolutionary Primatology (NYCEP), New York, New York

⁵Department of Anthropology, Baylor University, Waco, Texas

⁶Child and Brain Development Program, CIFAR, Toronto, Canada

⁷Department of Anthropology, University of Colorado-Colorado Springs, Colorado Springs, Colorado

⁸Department of Anthropology, Washington University in St. Louis, St. Louis, Missouri

⁹Department of Anthropology, Washington State University, Pullman, WA

¹⁰Department of Psychology, Boston College, Chestnut Hill, Massachusetts

¹¹Department of Anthropology, Yale University, New Haven, Connecticut

¹²Center for Global Health, University of Oregon, Eugene, Oregon

Correspondence

Alicia M. DeLouize, Department of Anthropology, 1218 University of Oregon, Eugene, OR 97403, USA.
Email: adelouiz@uoregon.edu

Funding information

Evonuk Foundation; Foundation for the National Institutes of Health, Grant/Award Number: DP1OD000516-04; Harvard University Sheldon Traveling Fellowship; Leakey Foundation; Lewis and Clark Exploration Grant; National Science Foundation (NSF) of the United States, Grant/Award Numbers: BCS-0824602, BCS-0925910, BCS-1340958, BCS-1341165, BCS-1650674, DGE-1144152, DGE-2011109300, SMA-1606852; Ryoichi Sasakawa Young Leaders Fellowship Fund; University of Oregon; University of Oregon Anthropology Department; University of Oregon Center for Latino/a and Latin American Studies; University of Oregon Institute of Cognitive and Decision Sciences; Wenner-Gren

Abstract

Objective: Anemia is an important global health challenge. We investigate anemia prevalence among Indigenous Shuar of Ecuador to expand our understanding of population-level variation, and to test hypotheses about how anemia variation is related to age, sex, and market integration.

Methods: Hemoglobin levels were measured in a total sample of 1650 Shuar participants (ages 6 months to 86 years) from 46 communities between 2008 and 2017 to compare anemia prevalence across regions characterized by different levels of market integration.

Results: Shuar anemia rates among children under 15 years (12.2%), adult women (10.5%), and adult men (5.3%) were less than half of those previously documented in other neo-tropical Indigenous populations. Anemia prevalence did not vary between more traditional and market integrated communities ($OR = 0.47$, $p = .52$). However, anemia was negatively associated with body mass index ($OR = 0.47$, $p = .002$).

Conclusions: Compared to other South American Indigenous populations, anemia prevalence is relatively low among Shuar of Ecuador and invariant

Foundation, Grant/Award Numbers: 7970, 8476, 8749, 9231, Engaged Anthropology Grant

with market integration. Understanding this pattern can provide valuable insights into anemia prevention among at-risk populations.

1 | INTRODUCTION

Anemia is a blood disorder characterized by insufficient oxygen transport by hemoglobin (Hb) in erythrocytes and is commonly defined as Hb levels below age- and sex-specific cutoffs (WHO, 2011). It afflicts ~27% of people worldwide, 89% of whom live in low- and middle-income countries (Kassebaum, 2016). Anemia is associated with negative health outcomes including fatigue, headaches, poor growth, reproductive disorders, low blood pressure, shortness of breath, reduced infection resistance, and lower cognitive function (Kassebaum, 2016). Dietary iron is needed for Hb function, with iron requirements varying by age, sex, menstrual cycle status, and reproductive state. Pregnancy and growth increase iron requirements leading to the highest rates of anemia in young children and women (Kassebaum, 2016; Kim, Yetley, & Calvo, 1993). Since anemia is readily treatable by addressing the underlying causes, the WHO considers this health challenge a global priority (WHO, 2011).

Anemia prevalence among South American Indigenous neotropical populations range from 19% to 100% in women, 6% to 79% in men, and 39% to 98% in children (Table 1). These generally high rates are attributed to low dietary iron and vitamin intake, blood loss from intestinal macroparasites, infection with *H. pylori*, and inflammation from other infections (Kassebaum, 2016). Yet, it is unclear how lifestyle changes within these populations influence anemia rates. Data from Shuar of Amazonian Ecuador allow for an analysis of how market integration (MI: the degree of consumption from and production for a market economy; Lu, 2007) affects anemia rates (Liebert et al., 2013). Most Shuar live in two main regions of the Morona-Santiago province in Ecuador: the Upano Valley (UV), which has greater access to healthcare facilities and market centers (Liebert et al., 2013; Urlacher et al., 2016), and Cross-Cutucú (CC), which is more remote and generally associated with high rates of parasitic infection (Cepon-Robins et al., 2014; Gildner et al., 2016, 2020).

We hypothesize that anemia prevalence will vary according to MI status, with region serving as a proxy for MI. Specifically, we hypothesize that CC Shuar will have higher rates of anemia than UV Shuar. Further, we expect anemia prevalence will be associated with sex, age, and body mass index (BMI).

2 | METHOD

2.1 | Participants

Shuar are an Indigenous lowland neotropical population of ~60 000–110 000 individuals. The CC region is more geographically isolated than the UV region, leading to differences in MI, lifestyle, economics, and diet (Urlacher et al., 2016). Traditional cultigens (including manioc, plantains, sweet potatoes, and yams) remain dietary staples in both regions. Reliance on foraging, fishing, and hunting vary but are generally greater in the CC region (e.g., Liebert et al., 2013; Urlacher et al., 2016). Shuar exhibit high rates of stunting and infectious disease burden but with complex inter- and intra-regional variation in prevalence (e.g., Blackwell, Pryor, Pozo, Tiwia, & Sugiyama, 2009; Gildner et al., 2016, 2020; Urlacher et al., 2018). Helminth infection rates of whipworm and roundworm are high (Cepon-Robins et al., 2014; Gildner et al., 2016, 2020).

Data were collected across nine field seasons from August 2008 through November 2017, as a part of the Shuar Health and Life History Project (shuarproject.org). Participants included 1650 Shuar (746 men, 904 women [26 pregnant]) ranging in age from 6 months to 86 years ($M = 19.20$, $SD = 16.73$), including 1062 participants from 29 UV communities (464 men, 598 women), and 588 participants from 17 CC communities (282 men, 306 women). In instances of repeat Hb measures, only the visit with the most recent Hb measurement was used in the analysis; thus, only one measurement was included per individual. Adults provided informed consent to participate. Parental consent and child assent were obtained for children and adolescents <15 years old, the legal age of consent in Ecuador. The study was approved by the IRB of the University of Oregon, by community leaders upon community consensus, and by the *Federación Interprovincial de Centros Shuar*.

2.2 | Measures

2.2.1 | Predictor variables

Age, height, and weight data were collected as described in detail elsewhere (Liebert et al., 2013; Urlacher et al., 2016). BMI was calculated using standard methods (weight [kg]/height [m]²).

TABLE 1 Comparison of Shuar mean Hb values and anemia prevalence with other South American populations

Group	Children				Adults				References
	Anemia (%)	WHO significance	Age (yrs)	Anemia (%)	WHO significance	Age (yrs)	Men		
							Anemia (%)	WHO significance	
Aruak/Karibe (Brazil)	66	Severe	0.5–10						Mondini, Rodrigues, Gimeno, & Baruzzi, 2009
Suruí (Brazil)	81	Severe	0.5–10						Orellana, Coimbra Jr, Lourenço, & Santos, 2006
Terena (Brazil)	51	Severe	2–4						Morais, Alves, & Fagundes-Neto, 2005
Indigenous (Brazil)	51	Severe	0.5–5						Leite et al., 2013
Pueblo Tacana (Bolivia)	42	Moderate	0.5–10						Benefice, Monroy, Jiménez, & López, 2006
Tsimane (Bolivia)	39	Moderate	0.5–15	23	Moderate	19	Mild	15+	Blackwell, unpublished
Qom/Toba (Argentina)				28	Moderate			18+	Goetz & Valeggia, 2017
Indigenous (Colombia)				31	Moderate	6	Mild	18+	Castillo, Oliveros, & Mora, 2014
Quechua (Peru)	49	Moderate	0.5–12						Cabada et al., 2015
Indigenous (Peru)	51	Severe	0.5–11						Anticona and San Sebastian 2014
Bari (Venezuela)	54	Severe	1–18						Diez-Ewald et al., 1997
Piaroa (Venezuela)	98	Severe	1–10	88	Severe	79	Severe	11+	Nieves García-Casal et al., 2008
Warao (Venezuela)	61	Severe	7–14						Wilbert & Layrisse, 1980
Interior Alto Ocamo Yanomamo (Venezuela)	84	Severe	0.5–14	100	Severe	58	Severe	15+	Grenfell et al., 2008
Ocamo Yanomamo (Venezuela)	57	Severe	0.5–14	100	Severe	8	Mild	15+	Grenfell et al., 2008
Quechua (Ecuador)				50	Severe	47	Severe	15+	Pleasant, 2014
Secoya (Ecuador)				19	Mild	26	Moderate	15+	Pleasant, 2014
Cofán (Ecuador)				31	Moderate	29	Moderate	15+	Pleasant, 2014
Shuar (Ecuador)	12	Mild	0.5–14	11	Mild	5	Normal	15+	This study
Trans Cutucú Shuar (Ecuador)	12	Mild	0.5–14	10	Mild	6	Mild	15+	This study
Upano Valley Shuar (Ecuador)	13	Mild	0.5–14	11	Mild	5	Normal	15+	This study

TABLE 2 Shuar anemia prevalence and hemoglobin levels by WHO age/sex criteria

Shuar age class	<i>n</i>	Hb <i>M</i> (mg/dl)	Hb <i>SD</i> (mg/dl)	Mild anemia (%)	Moderate anemia (%)	Severe anemia (%)	Anemia prevalence (%)
Children (0–4 yrs)	127	11.75	1.03	15.7	4.7	0.0	20.4
Children (5–11 yrs)	653	12.61	1.00	6.1	4.7	0.0	10.9
Children (12–14 yrs)	164	13.30	1.44	6.7	3.7	0.6	11.0
Nonpregnant Women (≥15 yrs)	382	13.31	1.33	6.2	3.7	0.5	10.5
Pregnant women (>15 yrs)	26	12.10	1.06	7.7	3.8	0.0	11.5
Men (≥15 yrs)	265	15.08	1.36	3.4	1.5	0.4	5.3

Note: The hemoglobin (Hb) cutoff values for anemia, mild anemia, moderate anemia, severe anemia, and anemia cutoff values were determined by current World Health Organization guidelines (WHO, 2011).

2.2.2 | Hemoglobin

A finger prick with a sterile, disposable lancet was used to collect one drop (~10 µl) of whole capillary blood into a microcuvette and analyzed using a HemoCue Hb201+ analyzer (HemoCue AB, Cypress, CA). No daily smokers were in the sample; therefore, Hb levels were only adjusted for minor variation in altitude between communities (WHO, 2011).

2.2.3 | Anemia

Anemia prevalence was evaluated according to WHO (2011) standard cutoff levels for defined anemia groups: children under 5 years, 5–11 years, and 12–14 years; nonpregnant women; pregnant women; and men aged 15 years or older.

2.2.4 | Statistics

Binary logistic regressions with mixed effects were performed to predict anemia from region, sex, and BMI; each predictor was entered into individual models. For analyses between anemia and region and anemia and BMI, sex, age, pregnancy status, and year of data collection were controlled for by including WHO defined anemia groups (WHO, 2011) and year of data collection as random intercepts. For analyses between anemia and sex, juveniles and adults were analyzed separately, pregnancy was included as a fixed effect for adults, and community and year of data collection were included as random intercepts. A binary logistic generalized additive regression was performed to predict anemia from thin plate splines fit to age using *REML*, while controlling for community, year of data collection, sex, and pregnancy status. Analyses were conducted in R 3.6.1 with $\alpha = .05$. All statistical assumptions were met before analysis.

3 | RESULTS

Based on WHO (2011) criteria, anemia rates among most Shuar age- and sex-groups identified in this study are considered of mild public health significance (Table 2). Anemia prevalence for children under 15 years (12.2%), adult women (10.5%), and adult men (5.3%) were less than half that of the next lowest neo-tropical Indigenous comparator (Tsimane; Table 1). Analyses showed that anemia rates did not differ significantly by region ($OR = 0.47$, $p = .52$).

In participants under 15 years, anemia prevalence was higher in boys than girls (13.9% and 10.5%, respectively), but the difference was not significant ($OR = 0.41$, $p = .09$). However, controlling for pregnancy, adult women ≥15 years had higher rates of anemia (11.8%) than adult men (5.6%; $OR = 0.69$, $p = .01$). Anemia risk was significantly associated with age, with rates highest among the youngest and oldest participants ($edf = 4.73$, $p < .001$; Figure S1). Finally, participants with higher BMI levels had lower prevalence of anemia ($OR = 0.47$, $p = .002$).

4 | DISCUSSION

Anemia prevalence among Shuar is generally low compared to other South American populations and comparable to levels in high-income countries (Kassebaum, 2016). Furthermore, there was no difference in anemia prevalence between UV and CC Shuar. Yet, BMI was negatively associated with anemia, suggesting that lifestyle (e.g., diet) might influence anemia rates, despite the lack of regional differences (Liebert et al., 2013). The low rate of anemia is surprising given that Shuar have a relatively heavy parasitic and infectious disease burden (Cepon-Robins et al., 2014; Gildner et al., 2016, 2020), severe enough to cause growth faltering in children (Urlacher et al., 2018). However, parasitic species known

to contribute most heavily to anemia (e.g., hookworm; Kassebaum, 2016) do not appear to be common in this population (Cepon-Robins et al., 2014; Gildner et al., 2020).

The low rates of anemia among Shuar and lack of regional differences may be due to a variety of factors. Traditional Shuar diet includes iron-rich fish and hunted game, but consumption varies between and within communities. Common staples (e.g., plantains, sweet potatoes, yams) also contain some non-heme iron. Wheat and corn, which inhibit iron absorption and limit bioavailability, are rarely consumed (Liebert et al., 2013). In Ecuador, the National Food Nutrition Program (PANN, 2000) for children 6–24 months includes iron fortification, which reduces anemia (Lutter et al., 2008), yet implementation appears uneven across Shuar communities and Shuar have lower anemia prevalence than Ecuadorians in general (Sackey, Weigel, & Armijos, 2003). Low levels of malaria could also be a contributing factor as the UV is generally above the malarial zone and endemic malaria in CC was reduced through control efforts.

Anemia prevalence among the Shuar is comparable to Europeans, Canadians, and Australians, and lower than Ecuador overall and other South American Indigenous populations (Coa and Ochoa, 2009; Le, 2016; Sackey et al., 2003; Stevens et al., 2013). These findings suggest that subsistence-based lifestyles are not always associated with high rates of anemia. For example, CC and UV anemia rates were similar despite UV Shuar generally exhibiting higher MI, including access to healthcare infrastructure and market centers, and CC having greater reliance on fish and game (Liebert et al., 2013; Urlacher et al., 2016). Future research will consider the relative contributions of dietary factors, infection rates, and infrastructure as contributors to the relatively low anemia prevalence among Shuar. The low prevalence of anemia and similarity between regions are unexpected findings: understanding these results could identify key factors that can be targeted to reduce high anemia rates among at-risk populations and to maintain low anemia prevalence among the Shuar as integration into the market economy accelerates.

ACKNOWLEDGMENTS

We thank the Shuar for their participation and hospitality. We also thank the *Federación Interprovincial de Centros Shuar* (FICSH), former Dirigentes de Salud Oswaldo Mankash and Favio Chinkim, and our numerous colleagues, research assistants and friends who made this research possible including: Ruby Fried; Tiffany Gandolfo; Estella, Luzmila, Katy, and Charo Jempekat; Samantha, Utitai; Marcia Saramiento; Dona Berta and

Don Guimo; Nelly Sardi; Cesar Kayap; and Medardo Tunki.

AUTHOR CONTRIBUTIONS

Alicia DeLouize: Data curation; formal analysis; visualization; writing-original draft; writing-review & editing. **Melissa Liebert:** Data curation; formal analysis; funding acquisition; investigation; writing-original draft; writing-review & editing. **Felicia Madimenos:** Funding acquisition; investigation; writing-review & editing. **Samuel Urlacher:** Funding acquisition; investigation; writing-review & editing. **Joshua Schrock:** Funding acquisition; investigation; writing-review & editing. **Tara Cepon-Robins:** Funding acquisition; investigation; writing-review & editing. **Theresa Gildner:** Funding acquisition; investigation; writing-review & editing. **Aaron Blackwell:** Formal analysis; funding acquisition; investigation; visualization; writing-review & editing. **Christopher Harrington:** Investigation. **Dorsa Amir:** Investigation; writing-review & editing. **Richard Bribiescas:** Investigation; writing-review & editing. **James Snodgrass:** Funding acquisition; methodology; project administration; supervision; writing-review & editing. **Lawrence Sugiyama:** Conceptualization; funding acquisition; investigation; methodology; project administration; supervision; writing-original draft; writing-review & editing.


CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT


The data that support the findings of this study are available on request by emailing the form found at <https://www.shuarproject.org/data-sharing> to shuarproject@gmail.com. The data are not publicly available due to privacy or ethical restrictions.

ORCID

Alicia M. DeLouize  <https://orcid.org/0000-0002-9045-6428>


Melissa A. Liebert  <https://orcid.org/0000-0001-8013-6773>




Felicia C. Madimenos  <https://orcid.org/0000-0001-5442-232X>

Samuel S. Urlacher  <https://orcid.org/0000-0002-6489-4117>

Tara J. Cepon-Robins  <https://orcid.org/0000-0002-4508-8507>

Theresa E. Gildner  <https://orcid.org/0000-0001-7486-5208>

Aaron D. Blackwell  <https://orcid.org/0000-0002-5871-9865>

Dorsa Amir  <https://orcid.org/0000-0003-0255-0228>
 James Josh Snodgrass  <https://orcid.org/0000-0002-7021-777X>
 Lawrence S. Sugiyama  <https://orcid.org/0000-0003-1279-0006>

REFERENCES

- Anticona, C., & San Sebastian, M. (2014). Anemia and malnutrition in indigenous children and adolescents of the Peruvian Amazon in a context of lead exposure: A cross-sectional study. *Global Health Action*, 7(1), 22888. <https://doi.org/10.3402/gha.v7.22888>
- Benefice, E., Monroy, S. L., Jiménez, S., & López, R. (2006). Nutritional status of Amerindian children from the Beni River (lowland Bolivia) as related to environmental, maternal and dietary factors. *Public Health Nutrition*, 9(3), 327–335. <https://doi.org/10.1079/PHN2005852>
- Blackwell, A. D., Pryor, G., Pozo, J., Tiwia, W., & Sugiyama, L. S. (2009). Growth and market integration in Amazonia: A comparison of growth indicators between Shuar, Shiwiar, and nonindigenous school children. *American Journal of Human Biology*, 21(2), 161–171. <https://doi.org/10.1002/ajhb.20838>
- Cabada, M. M., Goodrich, M. R., Graham, B., Villanueva-Meyer, P. G., Deichsel, E. L., Lopez, M., ... White, A. C. (2015). Prevalence of intestinal helminths, anemia, and malnutrition in Paucartambo, Peru. *Pan American Journal of Public Health Revista Panamericana de Salud Pública*, 37(2), 69–75.
- Castillo, M., Oliveros, A., & Mora, A. (2014). Description of the health status of Colombian indigenous communities through basic laboratory test. *Journal of Life Sciences*, 8(9), 789–793.
- Cepon-Robins, T. J., Liebert, M. A., Gildner, T. E., Urlacher, S. S., Colehour, A. M., Snodgrass, J. J., Madimenos, F. C., & Sugiyama, L. S. (2014). Soil-transmitted helminth prevalence and infection intensity among geographically and economically distinct Shuar communities in the Ecuadorian Amazon. *Journal of Parasitology*, 100(5), 598–607.
- Coa, R., & Ochoa, L. H. (2009). *Bolivia Encuesta Nacional de Demografía y Salud: ENDSA-2008*. Calverton, Maryland: Ministerio de Salud y Deportes and Macro International. [https://dhsprogram.com/pubs/pdf/FR228/FR228\[08Feb2010\].pdf](https://dhsprogram.com/pubs/pdf/FR228/FR228[08Feb2010].pdf)
- Diez-Ewald, M., Torres-Guerra, E., Layrisse, M., Leets, I., Vizcaino, G., & Arteaga-Vizcaino, M. (1997). Prevalence of anemia, iron, folic acid and vitamin B12 deficiency in two Bari Indian communities from western Venezuela. *Investigacion Clinica*, 38(4), 191–201.
- Gildner, T. E., Cepon-Robins, T. J., Liebert, M. A., Urlacher, S. S., Madimenos, F. C., Snodgrass, J. J., & Sugiyama, L. S. (2016). Regional variation in *Ascaris lumbricoides* and *Trichuris trichiura* infections by age cohort and sex: Effects of market integration among the indigenous Shuar of Amazonian Ecuador. *Journal of Physiological Anthropology*, 35(1), 28.
- Gildner, T. E., Cepon-Robins, T. J., Liebert, M. A., Urlacher, S. S., Schrock, J. M., Harrington, C. J., Madimenos, F. C., Snodgrass, J. J., & Sugiyama, L. S. (2020). Market integration and soil-transmitted helminth infection among the Shuar of Amazonian Ecuador. *PLoS One*, 15(7), e0236924.
- Goetz Teddy, G., & Vallengia Claudia. (2017). The ecology of anemia: Anemia prevalence and correlated factors in adult indigenous women in Argentina. *American Journal of Human Biology*, 29(3), e22947. <http://dx.doi.org/10.1002/ajhb.22947>.
- Grenfell, P., Fanello, C. I., Magris, M., Goncalves, J., Metzger, W. G., Vivas-Martínez, S., Curtis, C., & Vivas, L. (2008). Anaemia and malaria in Yanomami communities with differing access to healthcare. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, 102(7), 645–652. <https://doi.org/10.1016/j.trstmh.2008.02.021>
- Kassebaum, N. J. (2016). The global burden of anemia. *Hematology/Oncology Clinics*, 30(2), 247–308.
- Kim, I., Yetley, E. A., & Calvo, M. S. (1993). Variations in iron-status measures during the menstrual cycle. *American Journal of Clinical Nutrition*, 58(5), 705–709. <https://doi.org/10.1093/ajcn/58.5.705>
- Le, C. H. H. (2016). The prevalence of anemia and moderate-severe anemia in the US population (NHANES 2003–2012). *PLoS One*, 11(11), e0166635. <https://doi.org/10.1371/journal.pone.0166635>
- Leite, M. S., Cardoso, A. M., Coimbra, C. E., Welch, J. R., Gugelmin, S. A., Cabral, P., ... Lúcia Escobar, A. (2013). Prevalence of anemia and associated factors among indigenous children in Brazil: Results from the first National Survey of indigenous People's health and nutrition. *Nutrition Journal*, 12(69), <https://doi.org/10.1186/1475-2891-12-69>
- Liebert, M. A., Snodgrass, J. J., Madimenos, F. C., Cepon, T. J., Blackwell, A. D., & Sugiyama, L. S. (2013). Implications of market integration for cardiovascular and metabolic health among an indigenous Amazonian Ecuadorian population. *Annals of Human Biology*, 40(3), 228–242.
- Lu, F. (2007). Integration into the market among indigenous peoples: A cross-cultural perspective from the Ecuadorian Amazon. *Current Anthropology*, 48(4), 593–602.
- Lutter, C. K., Rodríguez, A., Fuenmayor, G., Avila, L., Sempertegui, F., & Escobar, J. (2008). Growth and micronutrient status in children receiving a fortified complementary food. *The Journal of Nutrition*, 138(2), 379–388. <https://doi.org/10.1093/jn/138.2.379>
- Mondini, L., Rodrigues, D. A., Gimeno, S. G., & Baruzzi, R. G. (2009). Nutritional status and hemoglobin values of Aruak and Karibe Indian children: Upper Xingu, Central Brazil, 2001–2002. *Revista Brasileira de Epidemiologia*, 12, 469–477.
- Morais, M. B., Alves, G. M. D. S., & Fagundes-Neto, U. (2005). Estado nutricional de crianças índias terenas: evolução do peso e estatura e prevalência atual de anemia. *Jornal de Pediatria*, 81(5), 383–389.
- Nieves Garcia-Casal, M., Leets, I., Bracho, C., Hidalgo, M., Bastidas, G., Gomez, A., ... Pérez, H. (2008). Prevalence of anemia and deficiencies of iron, folic acid and vitamin B 12 in an indigenous community from the Venezuelan Amazon with a high incidence of malaria. *IÓrgano Oficial de la Sociedad Latinoamericana de Nutrición*, 58, 12–18.
- Orellana, J. D., Coimbra, C. E., Jr., Lourenço, A. E. P., & Santos, R. V. (2006). Nutritional status and anemia in Suruí Indian children, Brazilian Amazon. *Jornal De Pediatria*, 82(5), 383–388.
- Pleasant, E. (2014). *Food for thought: Impacts of market integration on diet and health in indigenous communities in the northern Ecuadorian Amazon*. <https://doi.org/10.17615/dd9z-v962>.



- Sackey, M. E., Weigel, M. M., & Armijos, R. X. (2003). Predictors and nutritional consequences of intestinal parasitic infections in rural Ecuadorian children. *Journal of Tropical Pediatrics*, *49* (1), 17–23.
- Stevens, G. A., Finucane, M. M., De-Regil, L. M., Paciorek, C. J., Flaxman, S. R., Branca, F., ... Ezzati, M. (2013). Global, regional, and national trends in haemoglobin concentration and prevalence of total and severe anaemia in children and pregnant and non-pregnant women for 1995–2011: A systematic analysis of population-representative data. *The Lancet Global Health*, *1*, e16–e25. [https://doi.org/10.1016/S2214-109X\(13\)70001-9](https://doi.org/10.1016/S2214-109X(13)70001-9)
- Urlacher, S. S., Liebert, M. A., Josh Snodgrass, J., Blackwell, A. D., Cepon-Robins, T. J., Gildner, T. E., Madimenos, F. C., Amir, D., Bribiescas, R. G., & Sugiyama, L. S. (2016). Heterogeneous effects of market integration on sub-adult body size and nutritional status among the Shuar of Amazonian Ecuador. *Annals of Human Biology*, *43*(4), 316–329. <https://doi.org/10.1080/03014460.2016.1192219>
- Urlacher, S. S., Ellison, P. T., Sugiyama, L. S., Pontzer, H., Eick, G., Liebert, M. A., ... Snodgrass, J. J. (2018). Tradeoffs between immune function and childhood growth among Amazonian forager-horticulturalists. *Proceedings of the National Academy of Sciences USA*, *115*(17), E3914–E3921.
- WHO - World Health Organization (2011). *Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity*. Vitamin and Mineral Nutrition Information System. Geneva. Retrieved from <http://www.who.int/vmnis/indicators/haemoglobin>. Accessed November 2019.
- Wilbert, J., & Layrisse, M. (1980). . *Demographic and biological studies of the Warao Indians*, UCLA Latin American Studies Series, vol. 45. Los Angeles, CA: UCLA Latin American Center Publications.

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of this article.

How to cite this article: DeLouize AM, Liebert MA, Madimenos FC, et al. Low prevalence of anemia among Shuar communities of Amazonian Ecuador. *Am J Hum Biol.* 2021;e23590. <https://doi.org/10.1002/ajhb.23590>